



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : A61K 7/46, C11B 9/00, C11D 3/50	A1	(11) International Publication Number: WO 99/18926	
		(43) International Publication Date: 22 April 1999 (22.04.99)	
(21) International Application Number: PCT/GB98/03057	(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).		
(22) International Filing Date: 9 October 1998 (09.10.98)			
(30) Priority Data: 9721587.5 10 October 1997 (10.10.97) GB			
(71) Applicant (for all designated States except US): QUEST INTERNATIONAL B.V. (NL/NL); Huizerstraatweg 28, NL-1411 GP Naarden (NL).			
(72) Inventors; and (75) Inventors/Applicants (for US only): PERRING, Keith, Douglas [GB/GB]; 14 Malvern Road, Ashford, Kent TN24 8HS (GB). CLEMENTS, Christopher, Francis [GB/GB]; Inveresk, Lookers Lane, Saltwood, Hythe, Kent CT21 5HW (GB). MACMASTER, Angus, Peter [GB/GB]; The Barn, Robus Ceramics, Evington Park, Hastingleigh, Nr Ashford, Kent TN25 5JH (GB).	Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.		
(74) Agents: HARRISON, David, C. et al.; Mewburn Ellis, York House, 23 Kingsway, London WC2B 6HP (GB).			
(54) Title: PERFUME COMPOSITIONS			
(57) Abstract A perfume composition which gives good deposition and/or substantially improved deodorant effectiveness on textiles incorporating spandex® fibres. The perfume comprises a mixture of fragrance materials in which at least 60 % by weight of the composition comprises fragrance materials drawn from categories I and II.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LY	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

PERFUME COMPOSITIONS

This invention relates to perfumes, to laundry compositions containing such perfumes, and the use of these compositions to deposit perfume on
5 fabrics.

The use of perfumes in laundry products has been established for many years. Perfume is used to cover base odour and to provide fragrance notes which are attractive or pleasing to the consumer. Generally,
10 it is important that a perfume be able to perform well olfactively at a number of stages, for example, from product 'in the pack', during product use, on damp cloth after laundering and on dry cloth (i.e. after drying the damp cloth). Certain perfumes have the
15 ability to provide deodorant action against body odour, either when directly applied to human skin, or when included in a laundry product. Such perfumes are described in EP-B-3172, US-A-4304679, US-A-4278658, US-A-4134838, US-A-4288341 and US-A-4289641, US-A-
20 5482635 and US-A-5554588.

It is important that sufficient fragrance should be transferred onto the fabric to be perceptible after laundering or (if the perfume has deodorant properties) to yield the deodorant effect.

A number of techniques have been proposed for increasing perfume delivery (to) and/or perfume longevity (on) substrates such as skin, hair, fabric and hard surfaces. This includes the use of fixative materials in the perfume to depress perfume ingredient partial pressures (eg GB 1534231) thereby reducing evaporative loss, and the use of carriers (eg EP 332259) or microcapsules (eg EP 376385) to deliver perfume to fabric. These technologies may increase perfume presence on dry cloth but involve further processing steps and/or material costs.

At the present time, many garments are made from fabric which contains a mixture of fibres, a proportion of which are elastic, so that the fabric has the ability to stretch and to recover from stretch. Spandex fibres are commonly used for this purpose. The term "spandex" has been adopted as a generic term by the United States Federal Trade Commission to denote a manufactured fibre in which the fibre-forming substance is a long chain synthetic polymer composed of at least 85% of a segmented polyurethane. A discussion of such fibres can be found in "History of Spandex Elastomeric Fibres" by A.J. Ultee, which is a chapter starting at page 278 in *Made Fibres: Their Origin and Development*, edited by R.V. Seymour and R.S. Porter, Elsevier 1993. Spandex

fibres are also referred to as "elastane" or "elasthane" fibres.

Another discussion of such fibres is found under the heading "Segmented Polyurethanes" at page 613 of *Handbook of Textile Fibres* by J. Gordon Cook, 5th Ed. Merrow Publishing Company 1984. Further description of elastanes and their applications can be found in "Synthesefasern: Grundlagen, Technologie, Verarbeitung und Anwendung", B von Falkei (editor), Verlag Chemie (1981). Commercially available elastanes are well known, in particular as sold under the name LYCRA®, a registered trade mark of DuPont de Nemours and Company. Patents relating to such fibres include US-A-5000899, US-A-5288779 and US-A-5362432.

Summary of the Invention

We have now discovered certain perfumes which give good deposition and/or substantially improved deodorant effectiveness on textiles incorporating spandex fibres.

Broadly, the present invention provides a perfume composition comprising a mixture of fragrance materials in which at least 60% by weight of the composition comprises fragrance materials drawn from the two categories below:

Category I

hydroxylic materials which are alcohols,

phenols or salicylates, with an octanol/water partition coefficient (P) whose common logarithm ($\log_{10}P$) is 2.5 or greater, and a gas chromatographic Kovats index (as determined on polydimethylsiloxane as non-polar stationary phase) lying within the range 1050 to 1600.

Category II

esters, ethers, ketones or aldehydes, with an octanol/water partition coefficient (P) whose common logarithm ($\log_{10}P$) is 2.5 or greater, and a gas chromatographic Kovats index (as determined on polydimethylsiloxane as non-polar stationary phase) lying within the range 1300 to 1600.

Particularly preferred are category I materials with a partition coefficient whose common logarithm is 3.0 or greater and a Kovats index of 1100 up to 1600, and category II materials which are ethers, esters, or ketones with a Kovats index of 1350 up to 1600, and possessing one or more rings in their molecular structures.

It is envisaged that the perfumes of this invention will be incorporated into a laundry or other

composition for treatment of fabrics. This may be a detergent composition or presoak composition for washing the fabrics or a softening composition for softening the washed fabrics during rinsing and drying.

We have also discovered that the perfume may be incorporated into a composition used for treatment of yarn or new fabric, to provide a perfume benefit on new garments.

The benefit from the perfume compositions may be good deposition or retention of fragrance materials on the fabric. We have observed good deposition of a range of fragrance materials, especially fragrance materials which are of mid-range volatility (i.e. intermediate between the volatile perfume materials used as "top-notes" and the materials of low volatility which are customarily used as base notes in perfumes). These materials of mid-range volatility are often not perceptible on other fabrics such as cotton, polyamide and polyester after washing and drying.

Preferably, the perfume is a deodorant perfume giving a Malodour Reduction Value on cotton of at least 0.25, preferably at least 0.5, in the Malodour Reduction Value test described below and which is generally as given in EP-A-147191 and corresponding US-A-4663068.

With such perfumes we have observed that there is an enhanced deodorant benefit when the fabrics incorporate spandex fibres, compared to conventional fabrics such as cotton, polyamide and polyester without spandex. This can be measured using the Malodour Reduction Value test, modified by varying the test fabric instead of varying the perfume.

The Malodour Reduction Value Test

In this test, the Malodour Reduction Value of a deodorant perfume is measured by assessing its effectiveness, when applied to fabric, in reducing body malodour when the fabric so treated is placed in contact with the axillae (armpits) of a panel of human subjects, and held there for a standard period of time. From subsequent olfactory evaluation by trained assessors, a Malodour Reduction Value can be calculated so giving a measure of the effectiveness as a deodorant of the perfume under test.

Stage 1 is preparation of the perfume treated fabric.

A fabric is selected for the test and cut into 20 cm x 20 cm squares. A control fabric is likewise cut into squares. Both fabrics are then washed in a front-loading drum-type washing machine with a standard unperfumed washing powder containing the following ingredients:

Ingredient	Parts by weight
Sodium dodecylbenzene sulphonate	9.0
C ₁₃₋₁₅ alcohol 7EO	4.0
Sodium tripolyphosphate	33.0
Alkaline sodium silicate	6.0
Sodium carboxymethyl cellulose	1.0
Magnesium silicate	1.0
Ethylenediamine tetraacetic acid	0.2
Sodium sulphate	15.0
Water	10.8

The washed pieces of fabric are then rinsed with cold water and finally dried. The fabric squares so obtained represent "untreated" fabric, that is fabric devoid of perfume, other deodorant materials, dressing and other water-soluble substances that subsequently might adversely affect the Malodour Reduction Value Test.

The untreated pieces of fabric are divided into two batches, one of which may receive no further washing treatment and then represents the control fabric in the test. The other batch of fabric pieces is re-washed in the washing machine with the same standard fabric washing powder to which has been added 0.2% by weight of the perfume under test. The perfume treated pieces of fabric are then rinsed with cold water and dried again. The fabric squares so obtained represent "test" fabric, that is fabric onto which the

test perfume has been delivered.

When the intention is to test perfume properties, the control and test fabrics are the same, e.g. polyester or cotton shirt fabric and the "untreated" fabric serves as control without further washing. To test deposition on different cloths, the test fabric can differ from the control fabric, and both may be washed with the perfumed washing powder.

Stage 2 is the carrying out of the test. A team of three Caucasian female assessors of age within the range of 20 to 40 years is selected for olfactory evaluation on the basis that each is able to rank correctly the odour levels of the series of standard aqueous solutions of isovaleric acid listed below, and each is able to assign a numerical score, corresponding to the odour intensity of one of these solutions, to the body malodour of a shirt insert after has been worn in the axillary region by a male subject for a standard period of time.

A panel of 40 human subjects for use in the test is assembled from Caucasian male subjects of age within the range of from 20 to 55 years. By screening, subjects are chosen who develop axillary body malodour

that is not unusually strong and who do not develop a stronger body malodour in one axilla compared with the other. Subjects who develop unusually strong body malodour, for example due to a diet including curry or garlic, are not selected for the panel.

For two weeks before the start of the test, the panel subjects are assigned an unperfumed, non-deodorant soap bar for exclusive use when washing and are denied the use of any other type of deodorant or antiperspirant. At the end of this period, the 40 subjects are randomly divided into two groups of 20.

The "test" and "control" fabric pieces are then tacked into 40 clean cotton or polyester-cotton shirts in the underarm region in such a manner that in 20 shirts, the control fabric pieces are attached inside the left underarm region, and the test fabric pieces are attached in the right underarm region. For the remaining 20 shirts, the placing of control and test pieces of fabric is reversed.

The shirts carrying the tacked-in fabric inserts are then worn by the 40 panel members for a period of 5 hours, during which time each panellist performs his normal work function without unnecessary

exercise.

After this five hour period, the shirts are removed and the inserts detached and placed in polyethylene pouches prior to assessment by the
5 trained panel of assessors.

The malodour intensity of each fabric insert is evaluated by all three assessors who, operating without knowledge of which inserts are "test" and which are "control" and, without knowing the scores
10 assigned by their fellow assessors, sniff each fabric piece and assign to it a score corresponding to the strength of the odour on a scale from 0 to 5, with 0 representing no odour and 5 representing very strong odour.

15 Standard aqueous solutions of isovaleric acid which correspond to each of the scores 1, 2, 3, 4 and 5 are provided for reference to assist the assessors in the malodour evaluation. These are shown below:

Score	Odour level	Concentration of aqueous isovaleric acid (ml/l)
20 0	No odour	0
1	Slight	0.013
2	Definite	0.053
3	Moderate	0.22
4	Strong	0.87
25 5	Very strong	3.57

11

The scores recorded by each assessor for each fabric piece are averaged. The average score of the "test" fabric pieces is deducted from the average score of the "untreated" control fabric pieces to give
5 a Malodour Reduction Value.

As a check that the selection of panel subjects is satisfactory for operation of the test, the average score with untreated fabric pieces should be between 2.5 and 3.0.

10 Preferred deodorant perfumes are those which have a Malodour Reduction Value of at least 0.50, or 0.70, or 1.00. The higher the minimum value, the more effective is the perfume as a deodorant as recorded by the assessors in the Malodour Reduction Value Test. It
15 has also been noted that consumers, who are not trained assessors, can detect by self-assessment a noticeable reduction in malodour on soiled fabric such as shirts and underclothes where the Malodour Reduction Value is at least 0.30, so the higher the
20 Malodour Reduction Value above this figure, the more noticeable is the deodorant effect.

Perfume Materials and Preferences

As mentioned above, the perfumes of this

invention must contain a number of fragrance materials specified by the presence of chemical structural groups, octanol/water partition coefficient (P) and Kovats index.

5 The octanol-water partition coefficient (or its common logarithm 'logP') is well known in the literature as an indicator of hydrophobicity and water solubility (see Hansch and Leo, *Chemical Reviews*, 526 to 616, (1971), 71; Hansch, Quinlan and Lawrence, 10 *J.Organic Chemistry*, 347 to 350 (1968), 33). Where such values are not available in the literature they may be measured directly, or approximately estimated using mathematical algorithms. Software providing such estimations are available commercially, for 15 example 'LogP' from Advanced Chemistry Design Inc.

A requirement for $\log_{10}P$ of 2.5 or more calls for materials which are somewhat hydrophobic.

Kovats indices are calculated from the retention time in a gas chromatographic measurement 20 referenced to the retention time for alkanes [see Kovats, *Helv.Chim.Acta* **41**, 1915 (1958)]. Indices based on the use of a non-polar stationary phase have been used in the perfumery industry for some years as

a descriptor relating to the molecular size and boiling point of ingredients. A review of Kovats indices in the perfume industry is given by T Shibamoto in "Capillary Gas Chromatography in Essential Oil Analysis", P Sandra and C Bicchi (editors), Huethig (1987), pages 259 to 274. A common non-polar phase which is suitable is 100% dimethyl polysiloxane, as supplied for example under a variety of tradenames such as HP-1 (Hewlett-Packard), CP Sil 5 CB (chrompack), OV-1 (Ohio Valley) and Rtx-1 (Restek).

The perfume materials fall into two sets referred to as categories I and II, differing in their minimum values of Kovats index.

Category I includes alcohols of general formula ROH where the hydroxyl group may be primary, secondary or tertiary, and the R group is an alkyl or alkenyl group, optionally branched or substituted, cyclic or acyclic, such that ROH has partition coefficient and Kovats properties as defined above. Typically this group comprises monofunctional alkyl or arylalkyl alcohols with molecular weight falling within the range 150 to 230.

Category I also includes phenols of general

formula ArOH, where the Ar group denotes a benzene ring which may be substituted with one or more alkyl or alkenyl groups, or with an ester grouping $-CO_2A$, where A is a hydrocarbon radical. As at the ortho position relative to the hydroxy group, the compound is a salicylate. ArOH has partition coefficient and Kovats index as defined above. Typically this group comprises monohydroxylic phenols with molecular weight falling within the range 150 to 210.

Ingredients which are particularly preferred are those with a partition coefficient of 1000 or more, i.e. $\log_{10}P$ of 3 or more, and a Kovats parameter of 1100 up to 1600.

Some examples of hydroxylic ingredients which fulfil the above criteria for category I are listed as a table below. Materials which are in the preferred sub-set are marked with an asterisk. Semitrivial names are those used in standard texts known within the perfume industry, particularly: *Common Fragrance and Flavor Materials* by Bauer, Garbe and Surburg, VCH Publ., 2nd edition (1990), and *Perfume and Flavour Materials*, Steffen Arctander, published in two volumes by the author (1969).

Examples of fragrance materials in category I	
	1-(2'-tert-butylcyclohexyloxy)-butan-2-ol*
	3-methyl-5-(2',2',3'-trimethylcyclopent-3-enyl)-pentan-2-ol*
5	4-methyl-3-decen-5-ol*
	amyl salicylate*
	2-ethyl-4(2',2',3'-trimethylcyclopent-3'-enyl)but-2-enol* (Bangalol, TM)
	borneol*
10	carvacrol*
	citronellol*
	9-decenol*
	dihydroeugenol*
	dihydrolinalol*
15	dihydromyrcenol
	dihydroterpineol *
	eugenol
	geraniol*
	hydroxycitronellal*
20	isoamyl salicylate*
	isobutyl salicylate*
	isoeugenol*
	linalol
	menthol*
25	nerolidol*
	nerol*
	para tert-butyl cyclohexanol*
	phenoxanol*
	terpineol
30	tetrahydrogeraniol*
	tetrahydrolinalol
	tetrahydromyrcenol
	thymol*
	2-methoxy-4-methylphenol (Ultravanil, TM)
35	(4-isopropylcyclohexyl)-methanol*

Category II is esters, ketones, nitriles,

aldehydes or ethers which have an octanol-water partition coefficient whose common logarithm ($\log_{10}P$) is at least 2.5, and a Kovats index of 1300 up to 1600 (non-polar phase).

5 Ingredients of Category II are of general formula RX, where X may be in a primary, secondary or tertiary position and is one of the following groups: -COA, -OA, -CO₂A, -CN or -CHO. R and A are hydrocarbon residues, cyclic or non-cyclic and optionally
10 substituted. In some forms of this invention, category II excludes any material with a free hydroxy group, so that where a hydroxyl group is present, the material should be considered only for Category I membership. Typically, the materials of Category II are
15 monofunctional compounds with molecular weights in the range 160 to 230.

 Ingredients which are particularly preferred are those with a Kovats parameter falling within the range 1350 up to 1600, and possessing a molecular
20 structure containing a ring, such as phenyl or cycloalkyl.

 A number of fragrance materials which fulfil the above criteria for category II are listed in the

table below. Materials which are in the preferred subset are marked with an asterisk.

Examples of fragrance materials in category II	
5	1-methyl-4-(4-methyl-3-pentenyl)-3-cyclohexene-1-carbaldehyde*
	1-(5',5'-dimethylcyclohexenyl)-pent-en-1-one*
	2-heptyl cyclopentanone*
	2-methyl-3-(4'-tert-butylphenyl)propanal
	2-methylundecanal
10	2-undecenal
	2,2-dimethyl-3-(4'-ethylphenyl)-propanal
	3-(4'-isopropylphenyl)-2-methylpropanal
	4-methyl-4-phenylpent-2-yl acetate*
	allyl cyclohexyl propionate*
15	allyl cyclohexyloxyacetate*
	amyl benzoate*
	methyl ethyl ketone trimers (Azarbre, TM)
	benzophenone*
	3-(4'-tert-butylphenyl)-propanal (Bourgeonal, TM)
20	caryophyllene*
	cis-jasmone*
	citral diethyl acetal
	citronellal diethyl acetal
	citronellyl acetate
25	phenylethyl butyl ether (Cressanther, TM)
	damascone, alpha-*
	damascone, beta-*
	damascone, delta-*
	decalactone, gamma-*
30	dihydro isojasmonate*
	dihydrojasmone*
	dihydroterpinyl acetate
	dimethyl anthranilate*
	diphenyl oxide*

	diphenylmethane*
	dodecanal
	dodecen-2-al
	dodecane nitrile
5	1-ethoxy-1-phenoxyethane (Efetaal, TM)
	3-(1'-ethoxyethoxy)-3,7-dimethylocta-1,6-diene (Elintaal Forte (TM))
	4-(4'-methylpent-3'-enyl)-cyclohex-3-enal (Empetaal, TM)
10	ethyl tricyclo[5.2.1.0-2,6-]decane-2-carboxylate*
	1-(7-isopropyl-5-methylbicyclo[2.2.2]oct-5-en-2-yl)-1-ethanone* (Felvinone, TM)
	allyl tricyclodecenyl ether* (Fleuroxene, TM)
	tricyclodecenyl propanoate* (Florocyclene, TM)
15	gamma-undecalactone*
	n-methyl-n-phenyl-2-methylbutanamide (Gardamide, TM)
	tricyclodecenyl isobutyrate* (Gardocyclene, TM)
	geranyl acetate
	hexyl benzoate*
20	ionone alpha*
	ionone beta*
	isobutyl cinnamate*
	isobutyl quinoline*
	isoeugenyl acetate*
25	2,2,7,7-tetramethyltricycloundecan-5-one* (Isolongifolanone, TM)
	tricyclodecenyl acetate* (Jasmacyclene, TM)
	2-hexylcyclopentanone (Jasmatone, TM)
	4-acetoxy-3-pentyltetrahydropyran* (Jasmopyrane, TM)
30	ethyl 2-hexylacetoacetate (Jessate, TM)
	8-isopropyl-6-methylbicyclo[2.2.2]oct-5-ene-2-carbaldehyde (Maceal, TM)
	methyl 4-isopropyl-1-methylbicyclo[2.2.2]oct-5-ene-2-carboxylate*
35	methyl cinnamate
	alpha iso methyl ionone*
	methyl naphthyl ketone*
	nerolin

	nonalactone gamma
	nopyl acetate*
	para tert-butyl cyclohexyl acetate
5	4-isopropyl-1-methyl-2-[1'-propenyl]-benzene* (Pelargene, TM)
	phenoxyethyl isobutyrate*
	phenylethyl isoamyl ether*
	phenylethyl isobutyrate*
	tricyclodecenyl pivalate* (Pivacyclene, TM)
10	phenylethyl pivalate* (Pivarose, TM)
	phenylacetaldehyde hexylene glycol acetal*
	2,4-dimethyl-4-phenyltetrahydrofuran (Rhubafuran, TM)
	rose acetone*
15	terpinyl acetate
	4-isopropyl-1-methyl-2-[1'-propenyl]-benzene (Verdoracine, TM)
	yara*
	(4-isopropylcyclohexadienyl)ethyl formate

20 Selection of a combination of fragrance materials to give a deodorant effect is explained in patents such as US-A-430679 referred to earlier. Further systems of selection are given in US-A-5482635 and US-A-5554588 also mentioned above.

25 Such selections can be carried out using materials with preferred values of partition coefficient and Kovats index as discussed above.

 The perfume compositions of this invention can deliver fragrance or, with appropriate perfume a
30 deodorant benefit, to a range of fabrics, but the

benefit is particularly pronounced on fabrics containing spandex fibres.

The polymer which is spun into spandex fibres is a segmented polyurethane, that is a copolymer incorporating polyurethane linkages. The polymer generally contains so-called soft (i.e. lower melting) segments which may be polyalkylene ethers or polyesters and so-called hard (i.e. higher melting) segments which are portions derived from the reaction of an isocyanate and a chain extender which is typically a diamine.

The soft segments may be poly(tetramethylene)ethers, possibly containing substituted tetramethylene glycol residues as described in US-A-5000899. Organic diisocyanates which may be used include conventional diisocyanates, such as diphenylmethane-4,4'-diisocyanate, also known as methylene-bis (4-phenylisocyanate) or "MDI", 2,4-tolylene diisocyanate, methylene-bis(4-cyclohexylisocyanate), isophorone diisocyanate, tetramethylene-p-xylylene diisocyanate, and the like. MDI is preferred.

Chain extenders used in producing the hard

segment of the fibres preferably include one or more of ethylenediamine (EDA), 1,3-propylenediamine, 1,4-cyclohexanediamine, hydrogenated m-phenylenediamine (HPMD), 2-methylpentamethylene diamine (MPMD) and 1,2-propylene diamine. More preferably, the chain extender is one or more of ethylenediamine, 1,3-propylenediamine, and 1,4-cyclohexanediamine, optionally mixed with HPMD, MPMD and/or 1,2-propylenediamine.

10 Spandex fibres with poly(tetramethylene)ethers as the soft segments are marketed by Dupont de Nemours International S.A. under the registered trade mark LYCRA® of Dupont de Nemours and Company.

15 Spandex fibres are generally mixed with other fibres such as cotton, polyamide, wool, polyester and acrylics and made into yarn which is then made into fabric. The contents of spandex fibres is usually in a range from 0.5% by weight of the yarn or fabric up to 20 50%, more usually from 1% to 30% by weight of the yarn or fabric.

A wide range of garments may contain spandex fibres in the fabric, including active sports wear,

intimate apparel, hosiery and a variety of ready to wear casual clothing.

Fabric treatment compositions

Perfume compositions of the invention may be
5 incorporated into fabric treatment products for use in washing, rinsing drying or other treatment of fabrics.
Such a product may be any of:

a detergent composition for fabric washing,

a pretreatment composition for application
10 to selected areas of a garment prior to washing,

a pretreatment composition used in the
soaking of entire garments prior to washing,

a rinse conditioner composition for
15 softening washed fabrics during a rinsing step,

an additive composition for use jointly with
any of the above,

a fabric conditioning article intended to be placed with fabrics during drying, or

a spray for application directly to dry garments.

5 Such products can take a variety of forms including powders, bars, sticks, tablets, mousses, gels, liquids, sprays, and also fabric conditioning sheets to be placed with fabrics in a tumble dryer. The amount of perfume in such products may lie in a
10 range from 0.1% to 10% by weight of thereof. The incorporation of perfume into products of these types is known, and existing techniques may be used for incorporating perfume for this invention. It may be possible to incorporate perfume directly into a
15 product, but another possibility is, to absorb the perfume on a carrier material and then admix the perfume-plus-carrier- mixture into the fabric treatment product. This approach may notably be used with a solid fabric treatment product and an inert
20 particulate carrier.

A detergent composition to be perfumed with a perfume composition according to this invention will

normally contain a deterative surfactant in an amount from 2% to 50%, preferably 5 to 40% by weight of the composition, and a detergency builder in an amount from 5% to 80% by weight of the composition. The balance of the composition, if any, may include various ingredients known for inclusion in fabric washing detergents, including bleaching materials. Surfactants may be one or more soap or non-soap anionic, nonionic, cationic, amphoteric or zwitterionic surfactants, or combinations of these. Preferred surfactants which can be used are soaps and synthetic non-soap anionic and nonionic compounds. Mixtures of surfactants, for example mixed anionic or mixed anionic and nonionic compounds, are frequently used in detergent compositions.

Detergency builders are materials which function to soften hard water by solubilisation or other removal of calcium and to a lesser extent magnesium salts responsible for water hardness. The commonest water soluble inorganic builder is sodium tripolyphosphate. A further water soluble inorganic builder compound is sodium carbonate which is generally used in conjunction with a seed crystal to

accelerate the precipitation of calcium carbonate.
Common insoluble inorganic detergency builders are
zeolites and layered silicates. Organic detergency-
builders such as sodium citrate and polyacrylate can
5 also be used.

Some detergent compositions, usually liquids,
are formulated to contain from 5 to 50 wt% surfactant
but little or no detergency builder.

Other ingredients which are customarily
10 included in a detergent composition, although not
necessarily all together, include alkaline silicate,
peroxygen or chlorine bleaches, soil release agents,
heavy metal sequestrants, anti-redeposition agents
such as sodium carboxymethyl cellulose, enzymes,
15 enzyme stabilisers, fabric softening agents including
softening clays, fluorescent brighteners, antifoam
agents or conversely foam boosters and filler such as
sodium sulphate.

Pretreatment compositions for soaking of
20 soiled fabrics prior to the main washing step may
contain 5 to 80 wt% by weight detergency builder with
little or no surfactant. Such compositions frequently
include enzymes.

The amount of perfume in a detergent composition or a presoak composition is likely to lie in a range from 0.1 to 2% by weight of the composition.

- 5 A fabric conditioning composition may contain from 1% to 40% by weight of a fabric conditioning agent which may be a fabric softening agent, but may contain higher levels in a very concentrated product. Fabric softening agents are frequently nonionic or
- 10 cationic organic compounds incorporating at least one alkyl, alkenyl or acyl group of 8 or more carbon atoms. These include, but are not limited to:
- (i) quaternary ammonium and imidazolinium compounds and corresponding tertiary amines
- 15 and imidazolines incorporating at least one, preferably two, C8 to C30 alkyl or alkenyl groups; also including alkyl groups containing, ether, ester, carbonate or amide linkages, ethoxylated derivatives and
- 20 analogues of such compounds and also including compounds with more than one tertiary or quaternary nitrogen atom,

- (ii) aliphatic alcohols, esters, amines or
carboxylic acids incorporating a C8 to C30
alkyl, alkenyl or acyl group, including
esters of sorbitan and of polyhydric
5 alcohols,
- (iii) silicones, mineral oils and polyols such as
polyethylene glycol.

A number of fabric conditioning compounds are
set out in US-A-4137180, and EP-A-239910.

- 10 Fabric conditioning compositions for addition
to a rinse liquid are frequently in the form of
aqueous dispersions of the conditioning agent. They
can also be made in the form of powders.

- The amount of perfume in such conditioning
15 liquids and powders is usually 0.1% to 2% by weight.
Preferred levels can vary depending on the
concentration of softening agent and requirements of
the market.

- The amount of perfume in very concentrated
20 fabric conditioners may lie in the broader range 0.1%
to 10% by weight, preferably 2% to 8% by weight.

A fabric conditioning sheet is intended to be placed with damp, rinsed, laundry in a tumble dryer. Such a product contains a fabric conditioner, which may be a nonionic compound as mentioned above, soap and/or fatty acid, and which melts at temperatures encountered in a tumble dryer. This is carried on a porous sheet. Silicone oil may be included. The amount of perfume incorporated in such a product is usually from 2% to 10% of the product and frequently from 2% or 4% to 7% or 8% by weight of the product.

Another form of product for the treatment of fabrics is a carrier liquid containing perfume and packaged in an applicator which delivers the composition as a spray. Such a spray may be marketed as a "refreshing spray" for garments. In such a product, the content of perfume will generally lie in a range from 0.1% to 10% by weight of the liquid composition.

A further possibility is that the perfume is used in the treatment of yarn, or in the "finishing" of new fabric. This is a step in the wet processing of fabrics to improve hand or surface appearance of fabric. The fabric will typically be treated in an

aqueous treatment bath containing fabric softener to deposit at a level of up to 3% by weight of the fabric. Perfume according to this invention may be included in the bath to deposit at a level of 0.001% to 1% by weight of the fabric.

Example 1

A mixture of perfume ingredients was prepared and added to an unperfumed, but otherwise conventional, laundry detergent powder, to provide a perfume concentration of 0.5% by weight.

The perfumed powder was used to wash test cloths which had not previously been treated with any perfume. These were either all cotton, or 95% cotton with 5% spandex. After washing, the cloths were rinsed and then line dried overnight.

The perfume was extracted from the dry cloths with organic solvent, and the content of the perfume ingredients in the solvent extracts was determined by gas chromatography. If the concentration of an ingredient extracted from the spandex containing cloth was greater than from the all-cotton cloth by a factor of 5 to 20, the result was coded as a medium enhancement (M). If the concentration was greater by 20

or more, it was coded high(H) and if less than 5 or not measurable, it was coded(L).

The results obtained were as follows:

	Ingredient	K*	logP**	Enhancement	Category
5	Boisambrene Forte	1714	5.5	M	-
	Benzyl acetone	1206	2.0	M	-
	Citronellol	1209	3.6	H	I
	2,6-Dimethyl-heptan-2-ol	975	2.9	L	-
10	Jasmacyclene	1394	2.9	H	II
	Methyl salicylate	1167	2.3	L	-
	2-Phenylethanol	1087	1.4	L	-
	Terpinyl acetate	1331	4.0	H	II
	Tetrahydrogeraniol	1180	3.6	H	I
15	Tetrahydrolinalol	1083	3.5	H	I
	Tonalid	1840	6.4	M	-
	Yara	1416	3.2	H	II

* Measured on OV1 phase using capillary gc

** Measured or estimated using 'logP' software from

20 ACD Inc.

Example 2

Two perfume compositions embodying this invention and a comparative composition contained perfume ingredients in the specified categories, as

25 follows:

<u>Perfume</u>	<u>Category I</u>	<u>Category II</u>	<u>Other</u>
A	35.1	46.6	18.3
B	41.8	43.8	14.4
C	27.6	29.0	43.4

5. These were used in the procedure of the
 Malodour Reduction Value test, as above, using test
 cloths which were 95% cotton 5% spandex. For the
 control, unperfumed washing powder was used to wash
 all-cotton test cloths. The following results were
 10 obtained:

	<u>Perfume A</u>	<u>Perfume B</u>	<u>Perfume C</u>
Average panel score:	1.04	1.29	1.57
Control panel score:	2.46	2.46	2.46
15 Malodour Reduction Value:	1.42	1.17	0.89
Malodour Reduction Value as % of control score:	57.7	47.4	36.1

CLAIMS

1. A perfume composition which is a mixture of
fragrance materials characterized by containing at
least 60 wt % of ingredients which are fragrance
5 materials selected from:
- Category I) hydroxylic materials which are
alcohols, phenols or salicylates, with an
octanol/water partition coefficient (P)
whose common logarithm ($\log_{10}P$) is 2.5 or
10 greater, and a gas chromatographic Kovats
index (as determined on polydimethylsiloxane
as non-polar silicone stationary phase)
lying within the range 1050 to 1600, and
- Category II) esters, ethers, nitriles,
15 ketones or aldehydes, with an octanol/water
partition coefficient (P) whose common
logarithm ($\log_{10}P$) is 2.5 or greater, and a
gas chromatographic Kovats index (as
determined on polydimethylsiloxane as non-
20 polar silicone stationary phase) lying
within the range 1300 to 1600.

2. A perfume composition according to claim 1 containing at least 20 wt % of ingredients in said Category I.

3. A perfume composition according to claim 1 containing at least 20 wt % of ingredients in said Category II.

4. A perfume composition according to claim 1 containing at least 33 wt % of ingredients in said Category I.

5. A perfume composition according to claim 1 containing at least 30 wt % of ingredients in said Category II.

6. A perfume composition according to any one of the preceding claims containing at least 20 wt % of ingredients in said Category I selected from the following list:

1-(2'-tert-butylcyclohexyloxy)-butan-2-ol
3-methyl-5-(2',2',3'-trimethylcyclopent-3-enyl)-pentan-2-ol
4-methyl-3-decen-5-ol
amyl salicylate
2-ethyl-4(2',2',3'-trimethylcyclopent-3'-enyl)-but-2-enol (Bangalol, TM)
borneol
carvacrol
citronellol

	9-decenol
	dihydroeugenol
	dihydrolinalol
	dihydromyrcenol
5	dihydroterpineol
	eugenol
	geraniol
	hydroxycitronellal
	isoamyl salicylate
10	isobutyl salicylate
	isoeugenol
	linalol
	menthol
	nerolidol
15	nerol
	para tert-butyl cyclohexanol
	phenoxanol
	terpineol
	tetrahydrogeraniol
20	tetrahydrolinalol
	tetrahydromyrcenol
	thymol
	2-methoxy-4-methylphenol (Ultravanyl, TM)
	(4-isopropylcyclohexyl)-methanol

- 25 7. A perfume composition according to any one of the preceding claims containing at least 20 wt % of ingredients in said Category II selected from those in the following list:

30	1-methyl-4-(4-methyl-3-pentenyl)-3-cyclohexene-1-carbaldehyde
	1-(5',5'-dimethylcyclohexenyl)-pent-en-1-one
	2-heptyl cyclopentanone

	2-methyl-3-(4'-tert-butylphenyl)propanal	
	2-methylundecanal	
	2-undecenal	
	2,2-dimethyl-3-(4'-ethylphenyl)-propanal	
5	3-(4'-isopropylphenyl)-2-methylpropanal	
	4-methyl-4-phenylpent-2-yl acetate	
	allyl cyclohexyl propionate	
	allyl cyclohexyloxyacetate	
	amyl benzoate	
10	methyl ethyl ketone trimers (Azarbre, TM)	
	benzophenone	
	3-(4'-tert-butylphenyl)-propanal (Bourgeonal, TM)	
	caryophyllene	
	cis-jasmone	
15	citral diethyl acetal	
	citronellal diethyl acetal	
	citronellyl acetate	
	phenylethyl butyl ether (Cressanther, TM)	
	damascone, alpha-	
20	damascone, beta-	
	damascone, delta-	
	decalactone, gamma-	
	dihydro isojasmonate	
	dihydrojasmone	
25	dihydroterpinyl acetate	
	dimethyl anthranilate	
	diphenyl oxide	
	diphenylmethane	
	dodecanal	
30	dodecen-2-al	
	dodecane nitrile	
	1-ethoxy-1-phenoxyethane (Efetaal, TM)	
	3-(1'-ethoxyethoxy)-3,7-dimethylocta-1,6-diene (Elintaal Forte (TM))	
35	4-(4'-methylpent-3'-enyl)-cyclohex-3-enal (Empetaal, TM)	
	ethyl tricyclo[5.2.1.0-2,6-]decane-2-carboxylate	

	1-(7-isopropyl-5-methylbicyclo[2.2.2]oct-5-en-2-yl)-1-ethanone (Felvinone, TM)
	allyl tricyclodecenyl ether (Fleuroxene, TM)
	tricyclodecenyl propanoate (Florocyclene, TM)
5	gamma-undecalactone
	n-methyl-n-phenyl-2-methylbutanamide (Gardamide, TM)
	tricyclodecenyl isobutyrate (Gardocyclene, TM)
	geranyl acetate
10	hexyl benzoate
	ionone alpha
	ionone beta
	isobutyl cinnamate
	isobutyl quinoline
15	isoeugenyl acetate
	2,2,7,7-tetramethyltricycloundecan-5-one (Isolongifolanone, TM)
	tricyclodecenyl acetate (Jasmacyclene, TM)
	2-hexylcyclopentanone (Jasmatone, TM)
20	4-acetoxy-3-pentyltetrahydropyran (Jasmopyrane, TM)
	ethyl 2-hexylacetoacetate (Jessate, TM)
	8-isopropyl-6-methylbicyclo[2.2.2]oct-5-ene-2-carbaldehyde (Maceal, TM)
25	methyl 4-isopropyl-1-methylbicyclo[2.2.2]oct-5-ene-2-carboxylate
	methyl cinnamate
	alpha iso methyl ionone
	methyl naphthyl ketone
	nerolin
30	nonalactone gamma
	nopyl acetate
	para tert-butyl cyclohexyl acetate
	4-isopropyl-1-methyl-2-[1'-propenyl]-benzene (Pelargene, TM)
35	phenoxyethyl isobutyrate
	phenylethyl isoamyl ether
	phenylethyl isobutyrate
	tricyclodecenyl pivalate (Pivacyclene, TM)

	phenylethyl pivalate (Pivarose, TM)
	phenylacetaldehyde hexylene glycol acetal
	2,4-dimethyl-4-phenyltetrahydrofuran (Rhubafuran, TM)
5	rose acetone
	terpinyl acetate
	4-isopropyl-1-methyl-2-[1'-propenyl]-benzene (Verdoracine, TM)
	yara
10	(4-isopropylcyclohexadienyl)ethyl formate

8. A perfume composition according to any one of the preceding claims which is deodorant perfume having a malodour reduction value of from 0.5 to 3.0 as measured by the Malodour Reduction Test herein,

15 carried out using 100% cotton fabric for both the test and the control pieces.

9. Yarn or fabric containing spandex fibres, having a perfume composition according to any one of the preceding claims deposited on the said yarn or
20 fabric.

10. Use of a perfume composition as defined in any one of claims 1 to 8 in the treatment of yarn or fabrics containing spandex fibres.

11. A fabric treatment composition incorporating
25 a perfume according to any one of the preceding claims.

12. A composition according to according to claim
11 which is a detergent composition for washing
fabrics, containing from 2 wt % to 50 wt % surfactant
and from 0.1 wt % to 5 wt % of the perfume
5 composition.

13. A composition according to claim 11 which is
a rinse conditioner composition comprising an aqueous
dispersion containing 1 wt % to 40 wt % of a fabric
softening agent and from 0.1 wt % to 10 wt % of the
10 perfume composition.

14. A method of treating yarn or unworn fabric
comprising treating the yarn or fabric with a fabric
finishing composition, characterised by incorporating
a perfume according to any one of claims 1 to 8 into
15 the finishing composition.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 98/03057

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K7/46 C11B9/00 C11D3/50

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C11B C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X A	US 5 500 138 A (D.R.BACON ET AL.) 19 March 1996 see claims	1,8-11, 14 2-7,12, 13
X A	WO 97 31097 A (PROCTER & GAMBLE) 28 August 1997 see claims; tables 1,2 see page "Perfume B" see page 36 "perfume D" see page 40 "Perfume J"	1,8-11, 14 2-7,12, 13
-/-		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"S" document member of the same patent family

Date of the actual completion of the international search

4 February 1999

Date of mailing of the international search report

19/02/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentstraan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx: 31 65 1 epo nl,
Fax: (+31-70) 340-3018

Authorized officer

Van Moer, A

INTERNATIONAL SEARCH REPORT

I International Application No

PCT/GB 98/03057

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 554 588 A (J.M.BEHAN ET AL.) 10 September 1996 cited in the application see column 2, line 44 - column 6, line 23; claims	1,8-11, 14
A	---	2-7,12, 13
X	US 5 482 635 A (J.M.BEHAN) 9 January 1996 cited in the application see claims	1,8-11, 14
A	see column 3, line 1 - column 6, line 50	2-7,12, 13
A	---	1-8
	DATABASE WPI Section Ch, Week 7811 Derwent Publications Ltd., London, GB; Class A97, AN 78-20668A XP002092224 & JP 53 012432 A (UENO M), 3 February 1978 see abstract	
A	---	1-8
	DATABASE WPI Section Ch, Week 9429 Derwent Publications Ltd., London, GB; Class D21, AN 94-239062 XP002092225 & JP 06 172781 A (SHISEIDO CO LTD) , 21 June 1994 see abstract	
A	---	1-8
	US 4 604 488 A (Y.FUJIKURA ET AL.) 5 August 1986 see examples 4,6	
A	---	1-14
	US 4 304 679 A (D.C.HOOPER ET AL.) 8 December 1981 cited in the application see claims; examples 3,4	
A	---	1-14
	EP 0 147 191 A (UNILEVER) 3 July 1985 cited in the application see claims see page 5, line 10-16 see page 18, line 11 - page 21, line 30	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 98/03057

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4304679 A		IE 47904 B	25-07-1984
		IN 149765 A	10-04-1982
		JP 1667132 C	29-05-1992
		JP 3021599 B	25-03-1991
		JP 54107910 A	24-08-1979
		NL 7900211 A	16-07-1979
		SE 438688 B	29-04-1985
		SE 7801713 A	13-07-1979
		ZA 7900127 A	27-08-1980
EP 147191 A	03-07-1985	AU 575100 B	21-07-1988
		AU 3688884 A	04-07-1985
		BR 8406656 A	22-10-1985
		CA 1253084 A	25-04-1989
		GB 2151669 A,B	24-07-1985
		GR 82517 A	08-04-1985
		JP 3057960 B	03-09-1991
		JP 60155298 A	15-08-1985
		JP 2706410 B	28-01-1998
		JP 6122895 A	06-05-1994
		PT 79733 B	24-11-1986
		US 4663068 A	05-05-1987

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 98/03057

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5500138 A	19-03-1996	AU 3832195 A BR 9509410 A CN 1169157 A CZ 9701164 A EP 0787176 A JP 10507793 T WO 9612785 A	15-05-1996 03-11-1998 31-12-1997 12-11-1997 06-08-1997 28-07-1998 02-05-1996
WO 9731097 A	28-08-1997	US 5668094 A	16-09-1997
US 5554588 A	10-09-1996	CA 2082281 A DE 69221087 D DE 69221087 T EP 0545556 A ES 2104850 T JP 5255689 A MX 9206423 A ZA 9208578 A	09-05-1993 04-09-1997 13-11-1997 09-06-1993 16-10-1997 05-10-1993 01-05-1993 06-05-1994
US 5482635 A	09-01-1996	CA 2019222 A DE 69008048 D DE 69008048 T EP 0404470 A ES 2055329 T JP 2746735 B JP 3054300 A US 5501805 A AU 6578394 A BR 9406622 A EP 0695166 A WO 9424999 A JP 2791610 B JP 8509489 T ZA 9402960 A	19-12-1990 19-05-1994 03-11-1994 27-12-1990 16-08-1994 06-05-1998 08-03-1991 26-03-1996 21-11-1994 30-01-1996 07-02-1996 10-11-1994 27-08-1998 08-10-1996 30-10-1995
US 4604488 A	05-08-1986	JP 1412208 C JP 59116242 A JP 62020173 B CH 657868 A DE 3346213 A FR 2539624 A	27-11-1987 05-07-1984 06-05-1987 30-09-1986 05-07-1984 27-07-1984
US 4304679 A	08-12-1981	AR 227130 A AT 15879 A AU 521125 B AU 4322479 A BR 7900156 A CA 1122902 A CA 1218018 B DE 2900721 A DE 2967667 A DK 66078 A, B EP 0003172 A FI 780440 A FR 2414551 A GB 2012302 A, B GR 66547 A HK 42384 A	30-09-1982 15-05-1987 18-03-1982 19-07-1979 14-08-1979 04-05-1982 17-02-1987 19-07-1979 12-11-1987 13-07-1979 25-07-1979 13-07-1979 10-08-1979 25-07-1979 26-03-1981 18-05-1984